

DUFAZAN PLUS

Composition:

Contains per ml suspension:

Levamisole HCl 30 mg

Oxyclozanide..... 60 mg

Description:

Dufazan Plus is a bright yellow suspension containing levamisole and oxyclozanide. This combination is a broad-spectrum anthelmintic as well as a fasciolicide, active against almost all mature and immature stages of nematodes, lungworms and *Fasciola* spp in cattle, sheep and goat.

Indications:

Endoparasitic infections in cattle, sheep and goat, caused by parasites susceptible to levamisole or oxyclozanide. *Haemonchus* spp., *Trichostrongylus* spp., *Ostertagia* spp.*. (*effective against both immature and adult *Ostertagia* but not inhibited larvae of *Ostertagia*, *Haemonchus* and *Trichostrongylus axei*), *Cooperia* spp., *Nematodirus* spp., *Bunostomum* spp., *Oesophagostomum* spp., *Chabertia* spp., *Dictyocaulus* (lungworm), eyeworms (*Thelazia* spp.), *Fasciola* spp., *Paramphistomum* spp.

Contraindications:

Hypersensitivity to levamisole or oxyclozanide.

Dosage and administration:

For oral administration. Administration with drenching gun is recommended.

Recommended dose 1 ml per 4 kg body weight (ca. 7.5 mg Levamisole and 15 mg Oxyclozanide per kg of body weight)

For a correct dosage and to avoid sub-therapeutic doses, the body weight should be determined with accuracy

Side effects:

At the recommended dose levamisole rarely cause side effects. At higher doses transient symptoms may appear, these include head shaking, salivation and slight muscle tremors. These are more likely to occur in cattle than sheep. Oxyclozanide can cause slight softening of the faeces in cattle with sometimes increased frequency of defecation and transient inappetence.

Interactions:

Do not use within 14 days of organophosphorus compounds. Cattle may be treated simultaneously with phosmet, fenthion or metrifphonate fly dressings.

Precautions and warnings:

Dose accurately to prevent development of resistance caused by underdosing. Avoid skin contact. If skin contact occurs, wash immediately with soap and water. If eye contact occurs, flush eyes immediately with water and seek medical attention. Wash hands after use.

Withdrawal times:

Meat and offal: 28 days.

Do not use in animals producing milk for human consumption.

Storage conditions:

Store dark and below 25°C. Do not freeze.

Packing:

1 litre bottles.