

TRIM/SUL M 40/200 INJ

Composition:

Contains per ml solution:

Trimethoprim.....40 mg

Sulfamethoxazole200 mg

Description:

Trimethoprim is an antibacterial agent with a bacteriostatic activity against many gram-negative and gram-positive bacteria. Sulfamethoxazole is a sulfonamide antibiotic with bacteriostatic activity. The combination ('potentiated sulphas') is bactericidal against a wide variety of gram-positive and gram-negative bacteria as trimethoprim and sulfamethoxazole work synergistically.

Indications:

Infections in cattle, swine, sheep, goats and horses caused by microorganisms susceptible to the combined effects of trimethoprim and sulfamethoxazole, such as acute or chronic enteritis caused by *Salmonella choleraesuis* and some other *Salmonella* spp.; post-weaning diarrhoea caused by K88-positive, K99-positive or 987P *Escherichia coli*-strains; secondary bacterial infections with *Pasteurella multocida*, *Actinobacillus pleuropneumoniae*, *Streptococcus* spp and *Haemophilus parasuis*, during the course of viral disease.

Contraindications:

Hypersensitivity to trimethoprim or sulphonamides, blood dyscrasias or severe hepatic or renal impairment.

Dosage and administration:

For intramuscular (cattle, swine, sheep, goats) or slow intravenous (cattle, horse) administration.

Recommended dose 1 ml per 20 kg body weight twice daily for up to 5 days

To avoid under dosing the body weight should be determined with accuracy.

Side effects:

Transient injection site reactions. Anaemia, leucopenia and thrombocytopenia can be observed.

Interactions:

Do not administer concurrently with members of the Vitamin B complex, or local anaesthetics such as procaine and lidocaine or calcium based solutions.

Withdrawal times:

Meat: 14 days, Milk: 4 days

Storage conditions:

Store dark and below 25°C. Do not freeze.

After first opening it should be stored in dark and refrigerated conditions (2 – 8°C).

Packing: 100 ml vials.